

Wm C. White

admitted March 10th 1820

Pneumonia

1870

1871

the
first
was
the
day
the
to the
met
the
it is
money
the
other
is
to
the
only
of
of
2

1

Pneumonia, according to Haller, may be defined, an inflammation affecting either the viscera of the thorax or the membrane lining the cavity.

By some Nosologists, the morbid condition of the pulmonary organs is defined according to the part affected. The Pleura being inflamed, it is called Pleuritis or Pleurisy, when the parenchymata of the lungs are affected, it is termed Pneumonia Vera or Peripneumony, an engorged or suffocated state of the lungs is denominated Peripneumonia Notha or Bastard Peripneumony, a Rheumatic affection of the intercostal and contiguous ^{muscles} may be named Pleurodine or Bastard Pleurisy. But as it is my intention to treat only of the two former, it is useless to expatiate any longer on the great variety of cases, made out by nosologists.

As the symptoms and treatment of Pneumonia.

from
breast
of the
Pine
fell
down
The
thing
but
most
The
cots
tours
from
other
med
could
be p

monia vera and Pleuritis are much the same, and as I am not disposed to disagree from the most accurate observers, I shall treat of them both under the common head of Pneumonia.

Pneumonia may be distinguished by the following symptoms, Pyrexia, dolor in quadam thoracis parte, dyspnoea et tussis.

The generality of writers seem inclined to think the pain most frequent in the right side, but Dr Chapman is of opinion, that it is most frequently situated in the left.

The disease generally comes on with a cold stage and is attended by other symptoms of Pyrexia. Sometimes the Pyrexia is from the beginning accompanied with other symptoms, but frequently it is preceded before the other symptoms become considerable, and especially before the pain be felt. For the most part the pulse is

frequent, full, strong, hard, quick, and vibrating to the touch. But in a more advanced stage, it is sometimes weak and soft.

Bullen says, the difficulty of breathing is more considerable in inspiration, both because the lungs do not admit easily of dilatation, and because the dilatation aggravates the pain. The pain is generally greater, when the patient lies on the side affected.

The cough attending this disease is generally at first dry, but in a more advanced stage expectoration takes place and the matter spit up is various, both in consistence and colour, but frequently is streaked with blood.

The pain is not always confined to the sides, the sternum, the back between the shoulders &c are sometimes affected. When

sites
 to our
 affi
 onic
 still
 weigh
 time
 to the
 If
 it is
 dry is
 and
 cause
 prod
 whe
 inju
 sar a
 inf
 ten

situated in the side it is always confined to one spot. The place most frequently affected is about the 6 or 7th rib near the middle of its length. The pain is often very severe and pungent, but sometimes is more dull and obtuse, accompanied with a sense of weight. It is generally fixed, but sometimes it shoots from one part of the thorax to the other.

As regards the remote cause of the disease it may be ascribed to cold, applied to the body in any shape, so as to obstruct perspiration and determine to the lungs. There are other causes which may have a share in the production of the disease, such as any thing which shall obstruct, strain, or otherwise injure the Pneumonic organs. These causes are said to act more forcibly, when an inflammatory diathesis prevails in the system, and selects as its subjects, persons of a

noble
 by in
 the
 is bre
 cold,
 there
 life
 occur
 has a
 cipie
 in a,
 that
 H. w
 mot
 disce
 pure
 ates

robust make and vigorous health.

Pneumonia is said to occur most frequently in cold climates, and more particularly in the winter and springs seasons. As the disease is brought on by the vicissitudes of heat and cold, it may happen at any season in which these occur. It may occur at any period of life, but rarely before the age of puberty.

The Pneumonic inflammation frequently occurs as an epidemic, and from that cause has been ascribed by some authors* to a specific contagion, but I believe it to consist in a peculiar state of the atmosphere, and that it is brought into action by cold applied &c. which I have before considered as the remote causes.

Pneumonia like other inflammatory diseases, may terminate in resolution, suppuration or gangrene, but it is also terminated by an end peculiar to itself viz. an effu-

* Morgagne

in
large
space
the
who
dead
ad, in
to
the
part
pass
the
time
Dr
thor
term
the
by
to
act

sion of blood into the cellular texture of the lungs, which interrupting the circulation proves speedily fatal. "This (says Cullen) seems to be the most common termination of the disease when it is fatal; for upon dissecting persons dead of Pneumonic inflammation, it has appeared, that such an effusion had taken place".

From these dissections we also learn that there is an exudation which appears to consist partly of a soft viscid crust, often of a compact membranous form covering everywhere the pleura. This exudation appears also sometimes to be of a serous nature.

Dropsical effusions in the form of hydrothorax and Hydropericardii are another termination of the disease.

Another mode in which it terminates, is by the adhesion inflammation.

When the Pneumonic inflammation terminates by resolution, it is generally attended

by
V
H

at

gall

copi

the

dina

ian

mal

the

war

term

the

the

the

in

the

the

the

the

by some evident evacuation

The evacuation most commonly attending it, is the expectoration of a thick, white, or yellowish matter a little streaked with blood, copious and brought up without straining.

Other evacuations, are said to terminate the disease, such as epistaxis, bleeding from the hæmorrhoids, copious evacuations of a bilious matter by stool, and the voiding of urine, from which is deposited a copious sediment. A copious, warm, and fluid sweat is not an unfrequent termination of the disease.

But, prognostics are to be formed from observing the state of the principal symptoms.

'Tis to be feared the patient labours under violent inflammation, attended with difficult and painful respiration, when he can lie but on one side, when he can lie no other way than on his back, when he can breathe with tolerable ease in no other position than the

or
a
sho
the
the
dang
As
the
pae
A
imp
If
on
Sh
the
five
of
the
mo
pu

erect posture; when she is tormented with a violent cough attended with great pain; when the vessels of the neck become turgid, the face flushed, or the pulse irregular &c. These symptoms taken progressively mark the danger of the complaint.

As Pneumonia is most frequently resolved by expectoration, a hard, dry cough, attended with pain, indicates a violent disease.

A delirium coming on during pneumonic inflammation is always attended with danger.

If the disease prove fatal it is generally, on one of the days of the first week.

Should the disease be violent, and capable of resolution, it will generally happen during the first weeks, but in a more moderate state of the disease, it is frequently protracted till the second. As I have before mentioned, pneumonic inflammation sometimes ends in suppuration and gangrene.

at
a
be
a
with
with
eyes
of
on
cheap
now
eyes
how
him
pines
in
to
with
be
stre

The termination of Pneumonia in gangrene is a very rare occurrence; but if it should, it may be distinguished by the following symptoms, viz a relief from pain, the expectoration of purulent matter, streaked with dark coloured blood, foetid breath, rattling in the throat, a dejected countenance, dim eyes, a feeble, quick, and frequent pulse &c.

The treatment of this disease will be obvious on recurring to the symptoms. The remedy chiefly to be depended on is venisection, which should be used according to the urgency of the symptoms. Should we be called to a patient labouring under this complaint, we should bleed him until a relief of pain and freedom of expectoration be produced. If this cannot be done he should be bled till there is a disposition to syncope. In order, however, to do this with the greatest advantage, the venipuncture should be large enough to allow a bold and vigorous stream to flow; for nothing is better ascer-

same
Hedrick
cuddles
July
By
The
over
he
they
under
the
and
near
not
day
above
in
To
the
the

lained, in the practice of medicine, than the reduction of inflammation is more owing to the suddenness of the detraction, than to the quantity of blood drawn.

The disease seldom yields to one bleeding, however large the quantity taken away, for altho' the symptoms may be for the time relieved, they generally return with equally as much violence, and sometimes more. Should this be the case, we are again to recur to the lancet and so continue as long as the pain, cough, dyspnoea, and hardness of the pulse remain.

It is said by some practitioners, that it is not, incumbent to use the lancet after the 5th day, but it was the maxim of Dr. Rush, to abstract blood as long as the least vestige of inflammatory symptoms remained.

To prevent syncope let the patient be placed in a horizontal position.

To assist bleeding in general, should the

of
is
he
the
D
ess
rom
gran
be
is
ay
in
is
Be
not
of
the
can
cher

symptoms be urgent, we should procure topical bleeding by means of of leeches and the cupping-glass and scarificator applied over the seat of pain.

Dr Chapman is of opinion, that in the advanced stage of the disease, the capillary vessels become affected, and that however great the quantities of blood drawn by venesection, it will be of but little or no service, and that it is more topical bleeding is especially necessary. With the same reasons blisters may be employed. Their application should be as near as possible to the part affected.

But notwithstanding all this we should not be too hasty in laying aside the lancet. The pneumonic inflammation for several reasons, and among which we may advance the importance of the lungs to the animal economy, as well as the circulation of the whole mass of blood through them require

a-gr
xiny
del
relat
tom
singl
mend
drac
pres
stet
circu
nuso
he p
las
per
ing
F
ried
ty
comp

a greater degree of depletion, than that of any other venous; it is also a fact that women detract 20 30 or even 400g of blood in some instances, without producing the slightest symptom of syncope." This, says, Dr Chapman, is strikingly the case, that when bled in at the commencement of the disease, I have sometimes drawn as much as 4 or 500g of blood, without producing the slightest tendency to syncope.

After we are satisfied "continues he" that the circulation is sufficiently reduced, we should resort to local remedies" at the heat of which he places the vesicatory applications. There has been considerable controversy as to what period they should be applied; some considering that they should be applied at the onset of the disease, before venesection has been resorted to any extent, whilst others think them injurious if applied before the inflammatory symptoms are considerably reduced by blood-letting.

Do
ofin
a
plu
tind
the
off
gain
a
B
wne
blo
del
drea
wh
the
of
ten
of
lax

Dr Chapman is inclined to adopt the latter opinion.

As I have before mentioned, leeches, applied as near as possible to the seat of pain, tend considerably to lessen the violence of the complaint, and indeed so great is their efficacy, that we should recur to them a gain, and again, at any subsequent period of the disease.

There are some cases of inflammation, in which, after the abstraction of as much blood by venesection as we can with safety detract, there still continues to be a considerable degree of activity of pulse, and in which the pain is still violent; in these cases, the abstraction of 6 or 8 ℥ of blood, by means of the scarificator and cuppinglafs acts often like a charm. Should the bowels need opening we should prescribe some gentle laxatives, as meadial salts, olivine, or manna.

ally
for
cho-
most
of
large
D
reho
even
gloss
of
in an
in a
of
slig
three
person
by
male

As regards Diaphoretics, this are generally considered deleterious except in the forming stage of the disease. The one chiefly used is the *acupara* *acumibena*. The usual mode of administering, is in the form of infusion, which may be drunk in as large a quantity as the stomach will bear.

During the progress of the disease, Venisections should rarely if ever be resorted to, and even then only such as are calculated to co-operate with bloodletting, in reducing the force of the circulation. Combinations of Ipecac. or antimony with nitre are usually preferred, and altho they are not given with a view of exciting nausea, if they should produce it slightly, they will operate by removing the stricture of the breast, and in all probability promote expectoration.

Expectoration being the means by which nature attempts to rid herself of this dis-

agree
to pre
are
tion
as a
second
rather
from
mole
mole
year
out
7
adapt
the
febr
two
sub
clog
large

agreeable complaint, we should endeavour to promote it by giving such medicines as are supposed to act by increasing the secretions from the glands of the throat and bronchiae. And to do this more effectually, we should second the effects of those medicines, by directing the inhalation of the steam arising from a warm infusion of some of the emollient herbs viz Marsh-mallows, chamomile flowers &c. with the addition of vinegar. These should be frequently inhaled throughout the day.

None of the expectorants are however well adapted to this disease. The gums and even the squills, by their irritation, increase the febrile symptoms, and render the cough tighter. When these inflammatory symptoms are subdued, however, and the bronchiae remain clogged with mucus, they are of ten, particularly the squills, of great service in promoting

exp
ppre
abbi
mo
mth
m
dore
vanc
m
2
ma
do
m
L
la
h
len
ion
opia
crea

expectoration, and relieving both the cough and oppression. Ammonia is also sometimes serviceable. It may be given alone; but the usual mode of administering it is in combination with other substances, as the squills, Laudanum &c. The Lac ammoniaci may be ^{given} in the dose of from ℥ij to ℥ij with considerable advantage, in cases where the bronchiae are much clogged with mucus.

As an expectorant a decoction of Seneka snake root may be usefully employed.

As regards the use of opiates in pneumonia Physicians have differed considerably. But I cannot do better than extract the following passage from Haller. "From it appears that in the beginning of the disease, and before bleeding and blistering have produced some remission of pain and of the difficulty of breathing opiates have a very bad effect, by their increasing the difficulty of breathing, and in-

[Faint, illegible handwriting in cursive script, likely from the 18th or 19th century. The text is written in a single column across the page.]

[Faint, illegible handwriting on the right edge of the page, continuing from the main text or as a separate note.]

inflammatory symptoms. But in a more advanced state of the disease, when the difficulty of breathing has abated, and when the urgent symptom is a cough, proving the chief cause of the continuance of the pain and the want of sleep, opiates may be employed with great advantage and safety."

It is sometimes the case, that after the violence of the disease has abated, there still remains a tightness across the chest, a dry cough, difficult and scanty expectoration. In such cases, Dr. Chapman is in the habit of giving a combination of opium, Speecac and colomel, at stated periods, "and even (says he) should slight pyrexia be produced, it is no objection, but on the contrary exterminates more completely every vestige of the remaining disease."

Throughout the whole progress of the disease, the antiphlogistic regimen is to be strictly adhered to. -

